

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Current and future CO₂ emissions from drained peatlands in Southeast Asia

A. Hooijer¹, S. Page², J. G. Canadell³, M. Silvius⁴, J. Kwadijk¹, H. Wösten⁵, and J. Jauhiainen⁶

¹Deltares – Delft Hydraulics, P.O. Box 177, 2600 MH Delft, The Netherlands

²Department of Geography, University of Leicester, University Road, Leicester, LE1 7RH, UK

³Global Carbon Project, CSIRO Marine and Atmospheric Research, GPO Box 3023, Canberra, ACT 2601, Australia

⁴Wetlands International, P.O. Box 471, 6700 AL Wageningen, The Netherlands

⁵Alterra, Wageningen University and Research Centre, P.O. Box 47, 6700 AA Wageningen, The Netherlands

⁶Department of Forest Ecology, University of Helsinki, 00014 Helsinki, Finland

Received: 20 June 2009 – Accepted: 23 June 2009 – Published: 20 July 2009

Correspondence to: A. Hooijer (aljosja.hooijer@deltares.nl)

Published by Copernicus Publications on behalf of the European Geosciences Union.

7207

Abstract

Forested tropical peatlands in Southeast Asia store at least 42 000 Million metric tonnes (Mt) of soil carbon. Human activity and climate change threatens the stability of this large pool which has been decreasing rapidly over the last few decades owing to deforestation, drainage and fire. In this paper we estimate the carbon dioxide (CO₂) emissions resulting from drainage of lowland tropical peatland for agricultural and forestry development which dominates the perturbation of the carbon balance in the region. Present and future emissions from drained peatlands are quantified using data on peatland extent and peat thickness, present and projected land use, water management practices and decomposition rates. Of the 27.1 Million hectares (Mha) of peatland in Southeast Asia, 12.9 Mha had been deforested and mostly drained by 2006. This latter area is increasing rapidly as a result of increasing land development pressures. Carbon dioxide (CO₂) emission caused by decomposition of drained peatlands was between 355 and 855 Mt y⁻¹ in 2006 of which 82% came from Indonesia, largely Sumatra and Kalimantan. At a global scale, CO₂ emission from peatland drainage in Southeast Asia is contributing the equivalent of 1.3 to 3.1% of current global CO₂ emissions from the combustion of fossil fuel. If current peatland development and management practices continue, these emissions are predicted to continue for decades. This warrants inclusion of tropical peatland CO₂ emissions in global greenhouse gas emission calculations and climate mitigation policies. Uncertainties in emission calculations are discussed and research needs for improved estimates are identified.

1 Introduction

Peat deposits consist of plant remains (about 10% by weight of peat) and water (90%), accumulated in waterlogged and usually acidic conditions over thousands of years. Peatlands are the result of a fine balance between hydrology, ecology and landscape morphology (Page et al., 1999). A change in any of these three components will lead

7208

inevitably to a change in the rate of peat accumulation.

Human intervention has major impacts on peatland hydrology through rapid transformation of landscape structure and function unless appropriate water management is implemented (Hooijer, 2005a; Wösten et al., 2006a).

5 Lowland peatlands in Southeast Asia cover 27.1 Million hectares (Mha) (Wetlands International, 2003, 2004; FAO, 2004) of which over 22.5 Mha are in Indonesia where they make up 12% of the land area and over 50% of the lowland area by most definitions. Peat thicknesses range from 0.5 to 20 m (Page et al., 2002), with at least 17% over 4 m deep in Indonesia. This yields an estimated carbon store in Southeast Asian
10 peatlands of at least 42 000 Mt assuming a carbon content of 60 kg m^{-3} (Page et al., 2002).

Forested peatlands in Southeast Asia are being deforested, drained and often burned for agricultural development (mainly oil palm and pulpwood plantations). Widespread illegal logging, particularly in Indonesia, has also resulted in peat drainage
15 through construction of logging canals which leads to increased risk of fire (Page et al., 2002; Aldhous, 2004; Langner and Siegert, 2009). Recently domestic and international interest in using palm oil as a source of biofuel has contributed to further deforestation and drainage of peat swamp forest, particularly in Indonesia and Malaysia (Hooijer et al., 2006; Stone, 2007).

20 All these land use activities have impacts on the net greenhouse gas (ghg) balance of peatlands which are dominated by five flux components: i) CO_2 uptake by vegetation, ii) CO_2 emissions from peat decomposition, iii) CO_2 and other emissions from fires, iv) exports of dissolved and particulate organic carbon, and v) smaller role of emissions of methane (CH_4) and possibly nitrous oxides (N_2O).

25 This paper focuses on one of these components, namely CO_2 emissions from peat decomposition. Development of agriculture and other human activities on peatland requires drainage. This leads to aerobic conditions and higher redox potentials that favour microbial activity and nitrogen mineralization in the peat profile above the water table (Ueda et al., 2000; Jali, 2004) resulting in enhanced CO_2 loss by peat decompo-

7209

sition (Fig. 1).

Carbon emissions from drained tropical peatlands (other than from fires) have received limited attention in analyses of emissions from land use, land use change and forestry (LULUCF) (Canadell et al., 2007; Gullison et al., 2007), and are overlooked in
5 ghg emission budgets as considered by the UN Framework Convention on Climate Change (UNFCCC) (IPCC, 2007). While the links between peatland development and CO_2 emission are relatively well established for temperate and boreal peatlands (Minkinen et al., 2008; Oleszczuk et al., 2008) there is relatively little information on CO_2 emission from drained peatlands in the tropics.

10 In this paper we present the first geographically comprehensive analysis of CO_2 emission from the decomposition of organic matter from drained peatlands in Southeast Asia with particular reference to lowland peatlands in Indonesia, Malaysia, Papua New Guinea and Brunei. The analysis is based on data for peatland area, thickness and carbon content, and on rates of deforestation and drainage. In addition, we
15 establish a relationship between water table depth and peat decomposition in order to estimate present and future CO_2 emissions in Southeast Asia. Finally we discuss key uncertainties and future research needs for improved emission estimates.

2 Methods

2.1 Data

20 In order to estimate current and future CO_2 emissions from drained peatlands, the following information was obtained: i) where and how thick the peatlands are, ii) where they are drained and iii) to what depth, iv) what further deforestation and drainage developments can be expected, v) how much CO_2 emission is caused by drainage, and vi) how much peat carbon is available for oxidation. The required information is
25 addressed step-by-step below (for further details on methodologies and data sources see Hooijer et al., 2006).

7210

2.1.1 Peatland distribution and thickness

A peatland distribution map (Fig. 2) was obtained for the Indonesian islands of Sumatra and Kalimantan (Wetlands International, 2003; 2004). For the remaining areas, the Digital Soil Map of the World from the Food and Agriculture Organization (FAO, 2004) was used to determine peat percentage in soil classes. Peat thickness data for Sumatra, Kalimantan and Papua (Indonesia) were obtained from Wetlands International (2003, 2004). Average peat thicknesses for Malaysia, Brunei and Papua New Guinea were estimated conservatively on the basis of thicknesses in Indonesia. For the purpose of this study, we excluded smaller peatland areas found in other Southeast Asian countries (Philippines, Thailand and Vietnam) which are less studied and represent only a small fraction of the total area and carbon stock. Peatlands over 300 m above sea level were also excluded for the same reasons.

2.1.2 Distribution of drained peatlands in the year 2000

Distribution of drained peatlands in the year 2000 was derived from the Global Land Cover 2000 map (Bartholomé and Belward, 2005) that is based on a classification of "SPOT-VEGETATION" satellite images that have 1 km resolution. Sixteen land cover categories were divided into four drainage classes: "certainly drained if peatland" (cropland, which includes plantations and other large agricultural areas), "probably drained if peatland" (mosaics of cropland and other land uses), "possibly drained if peatland" (shrubland and burnt areas) and "probably not drained" (natural vegetation). Cells were assigned accordingly to drainage classes (Table 1) by fraction of area. Areas of peatland within each drainage class are presented in Table 2, by Province (in Indonesia), State (in Malaysia) and Country (outside Indonesia and Malaysia).

7211

2.1.3 Groundwater depths for the drainage classes

Groundwater depths for the drainage classes, i.e. average groundwater depths as presented in Table 1, were estimated from published data (Armentano and Menges, 1986; Murayama and Bakar, 1996; Wösten and Ritzema, 2001; Jauhiainen et al., 2004; Hooijer, 2005b; Melling et al. 2005; Ali et al. 2006) and field observations by the authors.

2.1.4 History and future trends in peatland drainage

Historical information on peatland cover, and therefore of drained area, was obtained from changes in forest cover between 1985 (Global Forest Watch, 2002) and 2000 (Bartholomé and Belward, 2005) (Table 2). The deforestation rate in peatlands over this period was $1.3\% \text{ y}^{-1}$ for Indonesia, varying from $0.5\% \text{ y}^{-1}$ in Papua Province to $2.8\% \text{ y}^{-1}$ in East Kalimantan Province. Similar rates apply to the other countries in Southeast Asia included in this analysis (Hooijer et al., 2006). These historical peatland deforestation rates per Province were projected to future years, assuming a "business as usual" continuation of current developments. Changes in relative areas within deforested peatland of the drainage classes "cropland", "mosaic cropland+shrubland" and "shrubland", as the total deforested area increases, were projected using relationships derived from distribution of drainage classes in Indonesian Provinces in 2000, as a function of the deforested area (Hooijer et al., 2006) (Fig. 3).

2.1.5 Relationship between groundwater depth and CO₂ emission

A relation was derived from the results of two types of emission studies. The first type of study is gas emission monitoring in relation to water depth (Armentano and Menges, 1986; Murayama and Bakar, 1996; Jauhiainen et al., 2004; Melling et al., 2005; Ali et al., 2006). The second type of study is long term monitoring of peat subsidence in drained peatlands, combined with peat carbon content and bulk density analysis to factor out the contribution of compaction from the total subsidence rate; the

7212

remainder is attributed to CO₂ emission (as reviewed by Wösten et al., 1997; Wösten and Ritzema, 2001). The analysis yields the following relation (Fig. 4):

$$\text{CO}_2 \text{ emission (t ha}^{-1} \text{ y}^{-1}) = 91 * \text{Groundwater depth (m)} [R^2 = 0.71, n = 8]$$

Where CO₂ emission is expressed in t ha⁻¹ y⁻¹ and groundwater depth is the average depth of the water table below the peat surface, expressed in metres. This linear relation implies that every 10 cm water table drawdown will result in an increase in CO₂ emission value of 9.1 t CO₂ ha⁻¹ y⁻¹.

2.1.6 Carbon content

Carbon content of Southeast Asian peat was taken to be 60 kg m⁻³ (Kanapathy, 1976; Neuzil, 1997; Page et al., 2002) and this value was applied to all areas.

2.2 Calculations

Using the data and relationships described above, the CO₂ emission from all geographical units was calculated as follows:

$$\text{CO}_2 \text{ emission} = \text{LU_Area} * \text{D_Area} * \text{D_Depth} * \text{CO}_2\text{-1m [t/y]}$$

Where:

LU_Area = peatland area with specific land use [ha]

D_Area = drained area within peatland area with specific land use [fraction]

D_Depth = average groundwater depth in drained peatland area with specific land use [m]

CO₂-1m = CO₂ emission at an average groundwater depth of 1 m = 91 [t CO₂ ha⁻¹ y⁻¹]

Different groundwater depths were applied to land cover types i.e. drainage classes as presented in Table 1 (“cropland”, “mosaic cropland and shrubland”, and “shrubland”), and emission calculated for the total area of each class within each geographic

7213

unit. Peatland drained to 0.95 m on average (considered “most likely” in plantations and other large-scale “cropland” areas; Table 1) will emit 86 t CO₂ ha⁻¹ y⁻¹. Peatland drained to 0.6 m depth on average (typical in small-scale agricultural areas, i.e. “mosaic cropland and shrubland”) for 88% of the area will emit 48 t CO₂ ha⁻¹ y⁻¹. Peatland drained to 0.33 m over half of the area (considered likely for “shrubland”, i.e., recently deforested areas, and burnt and degraded agricultural areas) will emit 15 t CO₂ ha⁻¹ y⁻¹. Subsequently, “minimum”, “likely” and “maximum” emission rates for the land cover types were calculated by varying drained area and groundwater depth in each drainage class as presented in Table 1. Overall emissions were estimated by multiplying CO₂ emissions per hectare of each land use type by the total area of the corresponding drainage class,

3 Results and discussion

3.1 Carbon fluxes and climate mitigation

Land cover trends from 1985–2000 (Table 2), extended to 2000–2006, indicate that about 47% of peatlands in Southeast Asia, or 12.9 Mha, were deforested by 2006. Projected rates of land use change within deforested areas in Southeast Asia over the same period suggest that 17% of this land is now drained extensively for large-scale agriculture (drainage class “cropland”), 67% is affected by moderately intensive drainage for small-scale agriculture (“mosaic cropland and shrubland”), and 16% is unmanaged, degrading non-agricultural peatland (“shrubland”). This results in an estimated total drained peatland area for 2006 of 11.1 Mha (9.5–12.7 Mh).

Carbon dioxide emission from organic matter decomposition in drained peatlands in 2006 is estimated to be 632 Mt y⁻¹ (355–855 Mt y⁻¹). This corresponds to an overall range of emissions between 6 and 100 t CO₂ ha⁻¹ y⁻¹. At present, Indonesia is the single largest CO₂ emitter from drained peatlands, responsible for 82% of Southeast Asian emission in 2006. Within Indonesia, Sumatra is the largest emitter closely fol-

7214

lowed by Kalimantan.

If current rates and practices of peatland development and degradation continue, CO₂ emission is expected to peak at 745 Mt y⁻¹ in 2015, followed by a steady decline over subsequent decades as the remaining peat deposits become increasingly depleted (Fig. 5). By 2030, emission is projected to decline to a likely value of 514 Mt y⁻¹ if peatland drainage continues without mitigation, and decline further to 236 Mt y⁻¹ by 2070. Cumulative CO₂ emission to 2006 from all peatlands in Southeast Asia was estimated at 9700 Mt (5300–13 700 Mt) and projected to be 25 900 Mt (17 200–31 000 Mt) by 2030 and 37 300 Mt (28 900–39 900 Mt) by 2070.

These emissions, on a unit area basis and for the same groundwater depth, are far higher in the tropics than in temperate and boreal areas, because the rate of aerobic decomposition is strongly influenced by temperature. A recent review of emissions from drained bogs and fens in temperate climates gives a median value of 4.1 t CO₂ ha⁻¹ y⁻¹ for arable land on drained fen peat Oleszczuk et al. (2008). This flux is one order of magnitude less than the lower end value of the emission range reported for drained tropical peatlands in our study.

Although significant uncertainties remain it is likely that CO₂ emissions from decomposition of drained peatlands in Southeast Asia of 355 to 855 Mt y⁻¹ in 2006 are equivalent to 1.3 to 3.1% of the 28 Billion metric tonnes of CO₂ y⁻¹ of global fossil fuel emissions during the same period (Canadell et al., 2007).

In addition to the permanent flux of emissions caused by peat decomposition, incidental emissions caused by peatland fires are at least of similar magnitude and much higher during El Niño-years. Average fire emissions have been estimated to be at least 1,400 Mt y⁻¹ CO₂ for 1997–2006 (Hooijer et al., 2006) and 469±187 Mt y⁻¹ CO₂ for 2000–2006 (van Werf et al., 2008). For the 1997–98 El Niño alone, estimates of fire emissions range from 6197 Mt CO₂ for Indonesia (2970 Mt–9423 Mt CO₂; Page et al., 2002) to 2662±836 Mt CO₂ for Indonesia, Malaysia and Papua New Guinea (van der Werf et al., 2008).

If the various numbers for CO₂ emissions for peat decomposition and peatland fires

7215

are combined, the minimum total emission would be 637 Mt y⁻¹ CO₂ (over 2000–2006, excluding the 1997–1998 El Niño) and the maximum 2255 Mt y⁻¹ CO₂ (over 1997–2006). Even though these estimates cover quite a range they consistently show that emissions from both peat decomposition and fires are large contributing fluxes to the net carbon balance of tropical peatlands in Southeast Asia.

The large magnitude of the emissions makes conservation of remaining forested tropical peatlands, and rehabilitation of degraded ones, a significant opportunity for carbon emission reductions. The concentrated nature of these emissions, they are produced on less than 0.1% of the global land area, makes them potentially easier to manage than many other emissions caused by multiple types of land conversion. Improved water management planning for whole hydrological units (peat domes) is the basis for conservation of peat resources.

Conservation and rehabilitation become even more critical when we place carbon dynamics from tropical peatlands in a long-term context that includes climate change (in addition to land use change). An analysis of climate projections to 2100 shows that 7 of 11 models agree on decreased rainfall during the dry seasons in a number of peatland regions of Southeast Asia (Li et al., 2007), and 9 of the models agree on greater interannual variability in dry season rainfall. These changes are strongest and most consistent across models for southern Sumatra and Borneo, where most peatland in Indonesia occurs. Decreased rainfall during the dry season will result in lower water tables exposing larger carbon stocks to aerobic conditions and so enhancing decomposition and CO₂ emissions. Already multiple ENSO events since 1997 have shown the characteristics of predicted future climates for the region and the positive feedbacks on carbon emissions between low rainfall events and intense land use of peatlands involving deforestation and drainage (Ali et al., 2006).

A post-Kyoto treaty after 2012 which includes carbon credits from Reduced Emissions from Deforestation and Degradation (REDD) is one of the most important opportunities for tropical peatlands to be valued for their environmental importance. This development will largely determine the opportunities for improved management, reha-

7216

bilitation and conservation, and consequently the magnitude of ghg emissions in the future.

Management and conservation of tropical peatlands clearly expose the connectivity and complexities between local development agendas, and global agendas on climate change and conservation of wetlands and biodiversity. Synergistic opportunities exist for sustainable regional development and climate change mitigation through supporting peatland management practices that result in reduced carbon emissions and enhanced forest conservation.

3.1.1 Uncertainties and research needs

In this section we discuss the main uncertainties of this analysis and identify the new research needs to improve future estimates of carbon emissions from peat decomposition.

Peat thickness of many regions in Indonesia is not well known. Peat thicknesses tend to be greatest in the central parts of the inaccessible, dome-shaped peat bodies, which are often tens of kilometres across. Most of the measurements on peat thicknesses are nearer to the fringes. New carbon estimates will improve by acquiring new datasets on peatland thickness particularly for Malaysia and Papua New Guinea for which little data is available.

Data on *extent and distribution* of peatlands need to be improved, especially for regions outside of Kalimantan and Sumatra.

Carbon content of Southeast Asian peat was assumed to be 60 kg C m^{-3} in this and several earlier studies (Wösten et al., 2001; Page et al., 2002). Carbon contents between 50 kg C m^{-3} (Rieley et al., 2008) and up to 90 kg C m^{-3} (Wetlands International, 2003; 2004) have been reported. This reflects a significant spatial variation in peat carbon content which can only be improved with additional measurements of peat bulk density.

Drainage depth classification as derived from the GLC 2000 global land cover classification (Bartholomé and Belward, 2005) needs to improve with more drainage classes

7217

to encompass the diversity of land uses and drainage depths. For example, areas in Papua (Indonesia) are classified as “mosaic cropland+shrubland” while they are known to actually be savannah-like swamp landscapes created by traditional land management techniques requiring regular burning (Silvius and Taufik, 1990). These areas are generally not “drained” in the normal sense but agriculture often takes place on elevated islands of dug up organic mud (from the submerged swamp soil).

The *percentage of peatland drained* within drainage classes was conservatively estimated from field surveys which did not cover all peatland regions, and focussed on deforested peatland. The percentage of drained peatland may be considerably larger than assessed here, as several interventions in the hydrological system are not taken into account. These include drainage in forested areas for log transport canals where legal or illegal logging takes place, and through the impacts of plantation and roadside drainage which often extend over distances of kilometres into adjacent forested areas.

Estimated likely *groundwater depths* (i.e. drainage depths) used in this analysis (Table 1) are somewhat greater than depths recommended in existing management guidelines. However in the case of intensively drained croplands and plantations they are shallower than depths often observed by the authors in practice; drainage depths over 1 m are common in oil-palm and pulp wood plantations. There is a need for an extended monitor system of groundwater depths in a range of peatland types under different management.

Values for *2006 land use* were projected from GLC 2000 data for the year 2000, corrected for deforestation trends, as this is still the most up-to-date published and validated land use dataset available for all of Southeast Asia. Continued improvement and update of land use data are required.

Projections have not taken into account *peatland drainability and future management responses*. When subsidence brings the peat surface close to the drainage base, resulting in increased flooding and reduced agricultural productivity, they may be abandoned and drainage intensity would decline. In such cases CO_2 emissions may be reduced. Part of the carbon stock in peatlands is below the drainage base and may

never be oxidized. However, a common observation is that drainage systems in abandoned peatlands continue to draw down water levels for decades, because no funding is available for canal blocking.

The *relation between CO₂ emission and average groundwater depth* as affected by drainage is a very sensitive parameter in the calculations, and has proven difficult to establish. There are two sources of information: gas flux measurements and subsidence monitoring. Gas flux measurements can be difficult to interpret because CO₂ emission resulting from peat oxidation must be separated from plant root respiration. There is also a very limited dataset of gas flux measurements and even more limited dataset on annual fluxes over multiple years to capture the high interannual variability of the system. Determining net carbon loss by monitoring subsidence of the peat surface, as well as changes in peat bulk density and carbon content to distinguish the effects of peat oxidation from those of shrinkage, is a more feasible method for a large scale regional assessment as the one presented here. Subsidence measurements have the additional advantage that they account for lateral export of particulate and dissolved organic matter into rivers and canals, a flux that is missed in gas flux measurements. A much larger network of long-term subsidence measurements will be required to improve regional estimates and links to CO₂ emissions. New relationships need to be explored to best characterize the water table regimes; recent unpublished findings suggest that a relation between minimum water depth (e.g. the 25 percentile) and peat decomposition rate could be more appropriate.

We have used a linear *relationship between groundwater depth and CO₂ emission*, fitted through data points derived from 6 different studies (Fig. 4). This relation needs further development as more field data is acquired particularly on land management aspects, peat characteristics and the time since the start of drainage. Additional data will also allow exploring whether this relationship is in fact nonlinear. The linear relation used in this study is considered the best estimate currently available for groundwater depths between 0.5 and 1 m, which covers the range of the most common groundwater depths in the study region.

7219

The only form of carbon emission considered in this assessment is CO₂. *Methane* (CH₄) *emissions* from both undrained and drained peatlands are found to be modest in comparison with CO₂ (Jauhiainen et al., 2005, 2008; Rieley et al., 2008), but may still be significant from a climate perspective given that CH₄ is a much stronger greenhouse gas (23 times stronger in “CO₂ equivalents”). New continued CH₄ flux measurements over multiple years will confirm to what extent this gas plays a significant role in the net ghg balance of peatlands. Likewise, very limited information on nitrous oxides (N₂O) emissions in peatlands requires new continued measurements, particularly in agricultural areas with nitrogen inputs.

Acknowledgements. The work reported here has been possible with the support provided by a great number of people and organizations, especially: Rinus Vis, Marcel Ververs, Rolf van Buren, Marjolijn Haasnoot (Deltares | Delft Hydraulics), David Hilbert (CSIRO), Faizal Parish (Global Environment Centre), Fred Stolle (Global Forest Watch), Florian Siegert (Remote Sensing Solutions), Niels Wielaard (SarVision), Jack Rieley (University of Nottingham). The original assessment was funded from internal Deltares | Delft Hydraulics R&D sources, within the PEAT-CO₂ (Peatland CO₂ Emission Assessment Tool) research programme. The work was completed under the Singapore Delft Water Alliance (SDWA) Peatland Programme. This assessment is a contribution to a synthesis effort on the vulnerabilities of tropical peatlands carried out under the auspices of the Global Carbon Project, a joint project of the Earth System Science Partnership.

References

- Aldhous P: Borneo is burning, *Nature*, 432, 144–146, 2004.
- Ali, M., Taylor, D., and Inubushi, K.: Effects of environmental variations on CO₂ efflux from a tropical peatland in eastern Sumatra, *Wetlands*, 26, 612–618, 2006.
- Armentano, T. V. and Menges, E. S.: Patterns of change in the carbon balance of organic soil-wetlands of the temperate zone, *J. Ecol.*, 74, 755–774, 1986.
- Bartholomé, E. and Belward, A. S.: G. L. C. 2000: a new approach to global land cover mapping from Earth observation data, *Int. J. Remote Sens.*, 26, 1959–1977, 2005.

7220

- Canadell, J. G., Le Quéré, C., Raupach, M. R., Field, C. B., Buitenhuis, E. T., Ciais, P., Conway, T. J., Gillett, N. P., Houghton, R. A., and Marland, G.: Contributions to accelerating atmospheric CO₂ growth from economic activity, carbon intensity, and efficiency of natural sinks, *P. Natl. Acad. Sci.*, 104, 18866–18870, 2007.
- 5 FAO: Digital Soil map of the World, 2004.
- Global Forest Watch: The State of the Forest: Indonesia, Bogor, Indonesia: Forest Watch Indonesia, and Washington DC: Global Forest Watch, 2002.
- Gullison, R. E., Frumhoff, P., Canadell, J. G., Field, C. B., and Nepstad, D. C., Hayhoe, K., Avissar, R., Curran, L. M., Friedlingstein, P., Jones, C. D., and Nobre, C.: Tropical forests and climate policy, *Science*, 316, 985–986, doi:10.1126/science.1136163, 2007.
- 10 Hooijer, A.: Hydrology of Tropical Wetland Forests: Recent Research Results from Sarawak Peat Swamps, in: *Forests-Water-People in the Humid Tropics*, edited by Bonell, M. and Bruijnzeel, L. A., Cambridge University Press, 447–461, 2005a.
- Hooijer A.: Hydrological Assessment of Forest Plantation Impacts on Tropical Forested Peatlands; Kampar Peninsula, Sumatra, Indonesia, in: *Landscape-Level Assessment of Hydrological & Ecological Values in the Kampar Peninsular*, ProForest (UK) report to APRIL, 2005b.
- 15 Hooijer, A., Silvius, M., Wösten, H., and Page, S.: PEAT-CO₂, Assessment of CO₂ emissions from drained peatlands in SE Asia, Delft Hydraulics report Q3943/2006, 36 p., <http://peat-co2.deltares.nl>, 2006.
- 20 IPCC: Climate Change 2007: The Physical Scientific Basis, Summary for Policymakers, IPCC Secretariat, Geneva, 21 pp., 2007.
- Jali, D.: Nitrogen Mineralization in the Tropical Peat Swamps, in: *Proceedings of the 12th International Peat Congress*, Tampere 6.–11.6.2004, edited by Päivänen, J., 644–652, 2004.
- 25 Jauhiainen, J., Jaya, A., Inoue, T., Heikkinen, J., Martikainen, P., and Vasander, H.: Carbon Balance in Managed Tropical Peat in Central Kalimantan, in: *Proceedings of the 12th International Peat Congress*, Tampere 6.–11.6.2004, edited by: Päivänen, J., 653–659, 2004.
- Jauhiainen, J., Takahashi, H., Heikkinen, J. E. P., Martikainen, P. J., and Vasander, H.: Carbon fluxes from a tropical peat swamp forest floor, *Glob. Change Biol.*, 11, 1788–1797, 2005.
- 30 Kanapathy, K.: Fertilizer requirement on peat soils, *Malays. Agr. J.*, 50, 292–307, 1976.
- Langner, A. and Siegert, F.: Spatiotemporal fire occurrence in Borneo over a period of 10 years, *Glob. Change Biol.*, 15, 48–62, 2009.
- Li, W., Dickinson, R. E., Fu, R., Niu, G. Y., Yang, Z. L., and Canadell, J. G.: Future Precipitation

7221

- Changes and Their Implications for Tropical Peatlands, *Geophys. Res. Lett.*, 34, L01403, doi:10.1029/2006GL028364, 2007.
- Melling, L., Hatano, R., and Goh, K. J.: Soil CO₂ flux from three ecosystems in tropical peatland of Sarawak, Malaysia, *Tellus B*, 57, 1–11, 2005.
- 5 Minkinen, K., Byrne, K. A., and Trettin, C.: Climate impacts of peatland forestry, in: *Peatlands and Climate Change*, edited by Strack, M., International Peat Society, Jyväskylä, Finland, 98–122, 2008.
- Murayama, S. and Bakar, Z. A.: Decomposition of tropical peat soils, 2. Estimation of in situ decomposition by measurement of CO₂ flux, *JARQ-Jpn. Agr. Res. Q.*, 30, 153–158, 1996.
- 10 Neuzil, S. G.: Onset and Rate of Peat and Carbon Accumulation in Four Domed Ombrogenous Peat Deposits, Indonesia, in: *Biodiversity and Sustainability in Peatlands – Proceedings of the International Symposium on Tropical Peatlands*, Palangka Raya, Indonesia, edited by: Rieley, J. O. and Page, S. E., Samara Publishing, Cardigan, 55–72, 1997.
- Oleszczuk, R., Regina, K., Szajdak, L., Höper, H., and Maryganova, V: Impacts of Agricultural Utilization of Peat Soils on the Greenhouse Gas Balance, in: *Peatlands and Climate Change*, edited by Strack, M., International Peat Society, Jyväskylä, Finland, 70–97, 2008.
- 15 Page, S. E., Rieley, J. O., Shotyk, O. W., and Weiss, D.: Interdependence of peat and vegetation in a tropical peat swamp forest, *Phil. Trans. R. Soc. Lond. B*, 345, 1885–1897, 1999.
- Page, S. E., Siegert, F., Rieley, J. O., Boehm, H. V., Jaya, A., and Limin, S.: The amount of carbon released from peat and forest fires in Indonesia during 1997, *Nature*, 420, 61–65, 2002.
- 20 Rieley, J. O., Wüst, R. A. J., Jauhiainen, J., Page, S. E., Ritzema, H., Wösten, H., Hooijer, A., Siegert, F., Limin, S., Vasander, H., and Stahlhut, M.: Tropical Peatlands, Carbon stores, Carbon Gas Emissions and Contribution to Climate Change Processes, in: *Peatlands and Climate Change*, edited by: Strack, M., International Peat Society, Jyväskylä, Finland, 148–181, 2008.
- Silvius, M. J. and Taufik, A. W.: Conservation and Land Use of Kimaam Island. A survey report and compilation of existing information, PHPA/AWB Bogor, 1990.
- Silvius, M. J. and Giesen, W.: Integration of Conservation and Land-Use Development of Swamp Forest of East Sumatra, In: *Proceedings of the Workshop on Sumatra, Environment and Development: Its Past, Present and Future*, Bogor, 16.–18. September 1992, BIOTROP Special Publication No. 46, SEAMEO BIOTROP, Bogor, 1992.
- 30 Stone, R.: Can palm oil plantations come clean?, *Science*, 317, 1491, 2007.

7222

- Ueda, S., Go, C.-S. U., Yoshioka, T., Yoshida, N., Wada, E., Miyajima, T., Sugimoto, A., Boontanon, N., Vijarnsorn, P., and Boonprakub, S.: Dynamics of dissolved O₂, CO₂, CH₄, and N₂O in a tropical coastal swamp in southern Thailand, *Biogeochemistry*, 49, 191–215, 2000.
- Van der Werf, G. R., Dempewolf, J., Trigg, S. N., Randerson, J. T., Kasibhatla, P. S., Giglio, L., Murdiyarso, D., Peters, W., Morton, D. C., Collatz, G. J., Dolman, A. J., and DeFries, R. S.: Climate regulation of fire emissions and deforestation in equatorial Asia, *PNAS*, 105, 20350–20355, 2008.
- Wetlands International: Maps of peatland distribution and carbon content in Sumatera, 1990–2002, 2003.
- Wetlands International: Maps of peatland distribution and carbon content in Kalimantan, 2000–2002, 2004.
- Wösten, J. H. M., Ismail, A. B., and Van Wijk, A. L. M.: Peat subsidence and its practical implications: a case study in Malaysia, *Geoderma*, 78, 25–36, 1997.
- Wösten, J. H. M. and Ritzema, H. P.: Land and water management options for peatland development in Sarawak, Malaysia, *Int. Peat J.*, 11, 59–66, 2001.
- Wösten, J. H. M., Van den Berg, J., Van Eijk, P., Gevers, G. J. M., Giesen, W. B. J. T., Hooijer, A., Idris, A., Leenman, P. H., Satriadi Rais, D., Siderius, C., Silvius, M. J., Suryadiputra, N., and Wibisono, I. W.: Interrelationships between hydrology and ecology in fire degraded tropical peat swamp forests, *International Journal of Water Resources Development*, 22, 157–174, 2006.

7223

Table 1. CO₂ emission calculation steps and main parameters.

			Minimum	Likely	Maximum
1. Drained area (within land use class)	Large croplands, including plantations	%	100	100	100
	Mixed cropland/shrubland: small-scale agriculture	%	75	88	100
	Shrubland; recently cleared and burnt areas	%	25	50	75
2. Groundwater depth (within land use class)	Large croplands, including plantations	m	0.80	0.95	1.10
	Mixed cropland/shrubland; small-scale agriculture	m	0.40	0.60	0.80
	Shrubland; recently cleared and burnt areas	m	0.25	0.33	0.40
3. Relation between groundwater depth and CO ₂ emission: 91 t/ha/y CO ₂ emission per m depth.					
Unit CO ₂ emission (calculated from 1, 2 and 3)	Large croplands, including plantations	t ha ⁻¹ y ⁻¹	73	86	100
	Mixed cropland/shrubland: small-scale agriculture	t ha ⁻¹ y ⁻¹	27	48	73
	Shrubland; recently cleared and burnt areas	t ha ⁻¹ y ⁻¹	6	15	27

7224

Table 2. Lowland peatland distribution with land use and rate of forest cover loss.

GLC 2000 class:	Low land peatland area (km ²)	Shrubland + burnt			Mosaic: crop+shrub			Cropland	Forest cover 2000					Forest change	
		6	8	total	2	9	total	12	1	4	5	total		1985	85-'00
		Mosaics & Shrub Cover, shrub component dominant, mainly evergreen	Shrub cover, mainly deciduous, (Dry or burnt)	Total shrubland + burnt	Mosaic: Tree cover and Cropland (incl. very degraded and open tree cover)	Mosaics of Cropland / Other natural veg. (shifting cultivation in mountains)	Total mix cropland + shrub (small-scale agr.)	Cultivated and managed, non irrigated (mixed)	Tree cover, broadleaved, evergreen, closed and closed to open	Tree cover, regularly flooded, Mangrove	Tree cover, regularly flooded, Swampy	Total forest (including logged)		Global Forest Watch / World Res. Inst.	Annual change over the period 1985 - 2000
		% area	% area	% area	% area	% area	% area	% area	% area	% area	% area	% area	% area	% area	%/y
Total Indonesia	225234	4	2	7	3	24	27	5	27	4	30	61	81	-1.3	
Kalimantan	58379	15	4	20	2	17	19	3	30	2	27	58	87	-1.9	
Central Kalimantan	30951	19	2	22	2	15	18	3	33	1	24	57	90	-2.2	
East Kalimantan	6655	22	19	42	0	9	9	5	29	4	11	44	85	-2.8	
West Kalimantan	17569	5	1	7	2	17	19	1	28	3	43	74	92	-1.2	
South Kalimantan	3204	15	3	18	6	45	51	14	14	0	4	18	41	-1.6	
Sumatra	69317	0	1	1	3	34	37	10	14	2	35	52	78	-1.8	
D.I. Aceh	2613	0	0	0	4	28	32	8	37	0	22	59	87	-1.8	
North Sumatra	3467	0	2	2	3	39	42	20	20	1	16	36	76	-2.6	
Riau	38365	0	1	1	2	24	26	7	14	3	49	66	87	-1.4	
Jambi	7076	0	1	1	3	38	40	17	9	0	33	42	67	-1.7	
South Sumatra	14015	0	1	2	4	57	61	12	11	1	14	26	66	-2.6	
West Sumatra	2096	0	5	5	4	42	46	11	24	0	13	38	69	-2.1	
Papua	75543	0	1	2	4	20	25	1	36	9	27	72	80	-0.5	
Other Indonesia~	21995	4	2	7	3	24	27	5	27	4	30	61	81	-1.3	
Malaysia	20431	2	1	1	7	32	38	7	36	4	15	53	78*	-1.8*	
Peninsular	5990	0	1	1	4	47	50	13	37	0	0	37	78*	-2.8*	
Sabah	1718	8	2	10	3	28	31	17	21	21	2	43	86*	-2.9*	
Sarawak	12723	2	1	2	9	26	35	4	38	3	23	59	76*	-1.1*	
Brunei	646	3	1	4	1	9	10	2	39	6	39	84	85*	-0.2*	
Papua N. Guinea	25680	0	1	1	4	32	35	3	38	5	19	61	80*	-1.3*	
SE ASIA	271991	4	2	5	4	26	29	5	29	4	28	61	81*	-1.3*	

~ Land use distribution for 'Other Indonesia' assumed equal to Total Indonesia.

* 1985 forest cover outside Indonesia is estimated.




Fig. 1. Schematic illustration of CO₂ emission from drained peatlands.




Fig. 2. Forest cover on peatland in the year 2000. Note that FAO non-histosol soil classes with 20–40% peat are not shown, hence peat extent may be greater than shown – e.g. Papua New Guinea has significant peatland cover.

7227




Fig. 3. Trends and projections of land use change in lowland peatland in SE Asia.

7228




Fig. 4. Linear relation between groundwater depth in peatland and CO₂ emission caused by peat decomposition. The line has been fitted through published measurements in agricultural areas in peatland, including oil palm plantations. Measurements in forest and improductive degraded peatlands are excluded because these are not representative for agricultural areas. Measurements in sites where average water depth is reported to be within 0.3 m are also excluded, because such sites are not effectively drained and often subject to frequent inundation. Most measurements are gas flux measurements at the peat surface; the Wösten and Ritzema 2001 data point is based on analysis of subsidence records.

7229




Fig. 5. Historical, current and projected CO₂ emissions from peatlands, as a result of drainage (fires excluded). The increase in emissions is caused by progressive drainage of an increased peatland area. The following decrease is caused by peat deposits being depleted, starting with the shallowest peat deposits that represent the largest peatland area. The stepwise pattern of this decrease is explained by the discrete peat thickness data available (0.75 m, 1.5 m, 3 m, 6 m, 10 m).

7230