

Global Carbon Budget 2017

The work presented here has been possible thanks to the enormous observational and modelling efforts of the institutions and networks below

Atmospheric CO₂ datasets

NOAA/ESRL (Dlugokencky and Tans 2017)

Scripps (Keeling et al. 1976)

Fossil Fuels and Industry

CDIAC (Boden et al. 2017)

USGS, 2017

UNFCCC, 2017

BP, 2017

Consumption Emissions

Peters et al. 2011

GTAP (Narayanan et al. 2015)

Land-Use Change

Houghton and Nassikas 2017

Hansis et al. 2015

GFED4 (van der Werf et al. 2017)

FAO-FRA and FAOSTAT

HYDE (Klein Goldewijk et al. 2017)

LUH2 (Hurtt et al. 2011)

Atmospheric inversions

CarbonTracker Europe (van der Laan-Luijkx et al. 2017)

Jena CarboScope (Rödenbeck et al. 2003)

CAMS (Chevallier et al. 2005)

Land models

CABLE | CLASS-CTEM | CLM4.5(BGC) | DLEM | ISAM |

JSBACH | JULES | LPJ-GUESS | LPJ | LPX-Bern | OCN |

ORCHIDEE | ORCHIDEE-MICT | SDGVM | VISIT

CRU (Harris et al. 2014)

Ocean models

CCSM-BEC | CSIRO | MITgem-REcoM2 | MPIOM-

HAMOCC | MICOM-HAMOCC | NEMO-PISCES (CNRM) |

NEMO-PISCES(IPSL) | NEMO-PlankTOM5 | NorESM-OC

pCO₂-based ocean flux products

Jena CarboScope (Rödenbeck et al. 2014)

Landschützer et al. 2016

SOCATv5 (Bakker et al. 2016)

Full references provided in [Le Quéré et al 2017](#)

C Le Quéré UK | **RM Andrew** Norway | **GP Peters** Norway | **JG Canadell** Australia | **P Friedlingstein** UK | **R Jackson** USA | **S Sitch** UK | **JI Korsbakken** Norway | **J Pongratz** Germany | **AC Manning** UK

Thomas A. Boden USA | **Pieter P. Tans** USA | **Oliver D. Andrews** UK | **Vivek K. Arora** Canada | **Dorothee C. E. Bakker** UK | **Leticia Barbero** USA | **Meike Becker** Norway | **Richard A. Betts** UK | **Laurent Bopp** France | **Frédéric Chevallier** France | **Louise P. Chini** USA | **Philippe Ciais** France | **Catherine E. Cosca** USA | **Jessica Cross** USA | **Kim Currie** New Zealand | **Thomas Gasser** Austria | **Ian Harris** UK | **Judith Hauck** Germany | **Vanessa Haverd** Australia | **Richard A. Houghton** USA | **Christopher W. Hunt** USA | **George Hurtt** USA | **Tatiana Ilyina** Germany | **Atul K. Jain** USA | **Etsushi Kato** Japan | **Markus Kautz** Germany | **Ralph F. Keeling** USA | **Kees Klein Goldewijk** The Netherlands | **Arne Körtzinger** Germany | **Peter Landschützer** Germany | **Nathalie Lefèvre** France | **Andrew Lenton** Australia | **Sebastian Lienert** Switzerland | **Ivan Lima** USA | **Danica Lombardozzi** USA | **Galen McKinley** USA | **Nicolas Metz** France | **Frank Millero** USA | **Pedro M. S. Monteiro** South Africa | **David R. Munro** USA | **Julia E. M. S. Nabel** Germany | **Shin-ichiro Nakaoka** Japan | **Yukihiro Nojiri** Japan | **X. Antonio Padín** Spain | **Anna Peregon** France | **Benjamin Pfeil** Norway | **Denis Pierrot** USA | **Benjamin Poulter** USA | **Gregor Rehder** Germany | **Janet Reimer** USA | **Christian Rödenbeck** Germany | **Joyashree Roy** India | **Jörg Schwinger** Norway | **Roland Séférian** France | **Ingunn Skjelvan** Norway | **Benjamin D. Stocker** Spain | **Hanqin Tian** USA | **Bronte Tilbrook** Australia | **Ingrid T. van der Laan-Luijkx** The Netherlands | **Guido R. van der Werf** The Netherlands | **Libo Wu** China | **Steven van Heuven** The Netherlands | **Nicolas Viovy** France | **Nicolas Vuichard** France | **Anthony P. Walker** USA | **Andrew J. Watson** UK | **Andrew J. Wiltshire** UK | **Sönke Zaehle** Germany | **Dan Zhu** France

Atlas Team Members at LSCE, France

P Ciais | **A Peregon** | **P Peylin** | **P Brockmann** | **V Maigné** | **P Evano** | **C Nangini**

Communications Team

O Gaffney | **A Minns** | **A Scrutton**

Earth Syst. Sci. Data Discuss., <https://doi.org/10.5194/essd-2017-123>

Manuscript under review for journal Earth Syst. Sci. Data

This is just a preview and not the published paper.

© Author(s) 2017. CC BY 4.0 License.



Global Carbon Budget 2017

Corinne Le Quéré¹, Robbie M. Andrew², Pierre Friedlingstein³, Stephen Sitch⁴, Julia Pongratz⁵, Andrew C. Manning⁶, Jan Ivar Korsbakken⁷, Glen P. Peters², Josep G. Canadell¹, Robert B. Jackson⁸, Thomas A. Boden⁹, Pieter P. Tans¹⁰, Oliver D. Andrews¹, Vivek K. Arora¹¹, Dorothee C. E. Bakker⁶, Leticia Barbero^{12,13}, Meike Becker^{14,15}, Richard A. Betts^{16,4}, Laurent Bopp¹⁷, Frédéric Chevallier¹⁸, Louise P. Chini¹⁹, Philippe Ciais¹⁸, Catherine E. Cosca²⁰, Jessica Cross²⁰, Kim Currie²¹, Thomas Gasser²², Ian Harris²³, Judith Hauck²⁴, Vanessa Haverd²⁵, Richard A. Houghton²⁶, Christopher W. Hunt²⁷, George Hurtt¹⁹, Tatiana Ilyina⁵, Atul K. Jain²⁸, Etsushi Kato²⁹, Markus Kautz³⁰, Ralph F. Keeling³¹, Kees Klein Goldewijk³², Arne Körtzinger³³, Peter Landschützer³⁴, Nathalie Lefèvre³⁴, Andrew Lenton^{35,36}, Sebastian Liener^{37,38}, Ivan Lima³⁹, Danica Lombardozzi⁴⁰, Nicolas Metz³⁴, Frank Millero⁴¹, Pedro M. S. Monteiro⁴², David R. Munro⁴³, Julia E. M. S. Nabel⁵, Shin-ichiro Nakaoka⁴⁴, Yukihiro Nojiri⁴⁴, X. Antonio Padin⁴⁵, Anna Peregon¹⁸, Benjamin Pfeil^{14,15}, Denis Pierrot^{12,13}, Benjamin Poulter^{46,47}, Gregor Rehder⁴⁸, Janet Reimer⁴⁹, Christian Rödenbeck⁵⁰, Jörg Schwinger⁵¹, Roland Séférian⁵², Ingunn Skjelvan⁵¹, Benjamin D. Stocker⁵³, Hanqin Tian⁵⁴, Bronte Tilbrook^{35,36}, Ingrid T. van der Laan-Luijk⁵⁵, Guido R. van der Werf⁵⁶, Steven van Heuven⁵⁷, Nicolas Viovy¹⁸, Nicolas Vuichard¹⁸, Anthony P. Walker⁵⁸, Andrew J. Watson⁴, Andrew J. Wiltshire¹⁶, Sönke Zaehle⁵⁰, Dan Zhu¹⁸

<https://doi.org/10.5194/essd-2017-123>

comment

Towards real-time verification of CO₂ emissions

The Paris Agreement has increased the incentive to verify reported anthropogenic carbon dioxide emissions with independent Earth system observations. Reliable verification requires a step change in our understanding of carbon cycle variability.

Glen P. Peters, Corinne Le Quéré, Robbie M. Andrew, Josep G. Canadell, Pierre Friedlingstein, Tatiana Ilyina, Robert B. Jackson, Fortunat Joos, Jan Ivar Korsbakken, Galen A. McKinley, Stephen Sitch and Pieter Tans

Emissions of CO₂ from fossil fuels and industry did not change from 2014 to 2016, yet there was a record increase in CO₂ concentration in the atmosphere. This apparent inconsistency is explained by the response of the natural carbon cycle to the 2015–2016 El Niño event¹, but it raises important questions about our ability to detect a sustained change in emissions from the atmospheric record. High-accuracy calibrated atmospheric measurements, diverse satellite data, and integrative modelling approaches could, and ultimately must, provide independent evidence of the effectiveness of collective action to address climate change. This verification will only be possible if we can fully filter out the background variability in atmospheric CO₂ concentrations driven by natural processes, a challenge that still remains.

(0.2–3.8%) and in the rest of the world of 1.9% (0.3%–3.4%) (ref. ²). The increased fossil fuel and industry emissions technically bring an end to the three years of approximately constant emissions that persisted from 2014 to 2016. Land-use change emissions in 2017 should be similar to their 2016 level³, based on fire observations using satellite data. When combining CO₂ emissions from fossil fuels, industry, and land-use change, we project 2017 global emissions to be 41.5 ± 4.4 billion tonnes of CO₂, similar to 2015 levels. Even though the projected 2017 emissions match those of the record year in 2015, they are not expected to increase atmospheric CO₂ concentration as much as in 2015 because of reinvigorated carbon uptake in natural reservoirs after the 2015–2016 El Niño event (Fig. 1).




Fig. 1 Trends in CO₂ emissions and atmospheric CO₂ concentrations. Even though CO₂ emissions from fossil fuel and industry, and total emissions including land-use change, have been relatively flat from 2014 to 2016, atmospheric concentrations saw a record increase in 2015 and 2016 (hatched) due to El Niño conditions. We expect



EDITORIAL

Warning signs for stabilizing global CO₂ emissions

R B Jackson¹, C Le Quéré², R M Andrew³, J G Canadell⁴, G P Peters⁵, J Roy⁶ and L Wu⁶

RECEIVED
18 October 2017

ACCEPTED FOR PUBLICATION
27 October 2017

PUBLISHED
DD MM 2017

Original content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



E-mail: rob.jackson@stanford.edu


Abstract

Carbon dioxide (CO₂) emissions from fossil fuels and industry comprise ~90% of all CO₂ emissions from human activities. For the last three years, such emissions were stable, despite continuing growth in the global economy. Many positive trends contributed to this unique hiatus, including reduced coal use in China and elsewhere, continuing gains in energy efficiency, and a boom in low-carbon renewables such as wind and solar. However, the temporary hiatus appears to have ended in 2017. For 2017, we project emissions growth of 2.0% (range: 0.8%–2.9%) from 2016 levels (leap-year adjusted), reaching a record 36.8 ± 1.8 Gt CO₂. Economic projections suggest further emissions growth in 2018 is likely. Time is running out on our ability to keep global average temperature increases below 2 °C and, even more immediately, anything close to 1.5 °C.

<https://doi.org/10.1038/s41558-017-0013-9>

<https://doi.org/10.1088/1748-9326/aa9662>

GCP Website



More information, data sources and data files:
<http://www.globalcarbonproject.org/carbonbudget>
 Contact: c.lequere@uea.ac.uk

Global Carbon Atlas



More information, data sources and data files:
www.globalcarbonatlas.org
 (co-funded in part by BNP Paribas Foundation)
 Contact: philippe.ciais@lsce.ipsl.fr

All the data is shown in billion tonnes CO₂ (GtCO₂)

1 Gigatonne (Gt) = 1 billion tonnes = 1×10^{15} g = 1 Petagram (Pg)

1 kg carbon (C) = 3.664 kg carbon dioxide (CO₂)

1 GtC = 3.664 billion tonnes CO₂ = 3.664 GtCO₂

(Figures in units of GtC and GtCO₂ are available from <http://globalcarbonbudget.org/carbonbudget>)


Most figures in this presentation are available for download as PDF or PNG from tinyurl.com/GCB17figs along with the data required to produce them.

Disclaimer

The Global Carbon Budget and the information presented here are intended for those interested in learning about the carbon cycle, and how human activities are changing it. The information contained herein is provided as a public service, with the understanding that the Global Carbon Project team make no warranties, either expressed or implied, concerning the accuracy, completeness, reliability, or suitability of the information.

Anthropogenic perturbation of the global carbon cycle

Perturbation of the global carbon cycle caused by anthropogenic activities, averaged globally for the decade 2007–2016 (GtCO₂/yr)



The budget imbalance is the difference between the estimated emissions and sinks.


Source: [CDIAC](#); [NOAA-ESRL](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Fossil Fuel and Industry Emissions

Emissions from fossil fuel use and industry

Global emissions from fossil fuel and industry: 36.2 ± 2 GtCO₂ in 2016, 62% over 1990

● Projection for 2017: 36.8 ± 2 GtCO₂, 2.0% higher than 2016




Estimates for 2015 and 2016 are preliminary. Growth rate is adjusted for the leap year in 2016.

Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Top emitters: fossil fuels and industry (absolute)

The top four emitters in 2016 covered 59% of global emissions
China (28%), United States (15%), EU28 (10%), India (7%)




Bunker fuels are used for international transport is 3.1% of global emissions.

Statistical differences between the global estimates and sum of national totals are 0.6% of global emissions.

Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Emissions Projections for 2017


Global emissions from fossil fuels and industry are projected to rise by 2.0% in 2017
The global projection has a large uncertainty, ranging from +0.8% to +3.0%



Source: [CDIAC](#); [Jackson et al 2017](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Top emitters: fossil fuels and industry (per capita)


Countries have a broad range of per capita emissions reflecting their national circumstances



Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Top emitters: fossil fuels and industry (per dollar)

Emissions per unit economic output (emissions intensities) generally decline over time
China's intensity is declining rapidly, but is still much higher than the world average




GDP is measured in purchasing power parity (PPP) terms in 2010 US dollars.

Source: [CDIAC](#); [IEA 2016](#) GDP to 2014, [IMF 2017](#) growth rates to 2016; [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Top emitters: fossil fuels and industry (bar chart)


Emissions by country from 2000 to 2016, with growth rates indicated for the more recent period of 2011 to 2016



Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Alternative rankings of countries

Depending on perspective, the significance of individual countries changes.
Emissions from fossil fuels and industry.



GDP: Gross Domestic Product in Market Exchange Rates (MER) and Purchasing Power Parity (PPP)

Source: [CDIAC](#); [United Nations](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Fossil fuel and industry emissions growth

Emissions in the US, Russia and Brazil declined in 2016
Emissions in India and all other countries combined increased





Figure shows the top four countries contributing to emissions changes in 2016

Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Breakdown of global emissions by country


Emissions from OECD countries are about the same as in 1990
Emissions from non-OECD countries have increased rapidly in the last decade



Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Historical cumulative emissions by country

Cumulative emissions from fossil-fuel and industry were distributed (1870–2016):
USA 26%, EU28 22%, China 13%, Russia 7%, Japan 4% and India 3%




Cumulative emissions (1990–2016) were distributed China 20%, USA 20%, EU28 14%, Russia 6%, India 5%, Japan 4%
'All others' includes all other countries along with bunker fuels and statistical differences

Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Historical cumulative emissions by continent

Cumulative emissions from fossil-fuel and industry (1870–2016)

North America and Europe responsible for most cumulative emissions, but Asia growing fast




The figure excludes bunker fuels and statistical differences

Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Emissions from coal, oil, gas, cement


Share of global emissions in 2016:
coal (40%), oil (34%), gas (19%), cement (6%), flaring (1%, not shown)



Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Emissions by category


Emissions by category from 2000 to 2016, with growth rates indicated for the more recent period of 2011 to 2016



Source: [CDIAC](#); [Jackson et al 2017](#); [Global Carbon Budget 2017](#)

Energy consumption by energy type


Energy consumption by fuel source from 2000 to 2016, with growth rates indicated for the more recent period of 2011 to 2016



Source: [BP 2017](#); [Jackson et al 2017](#); [Global Carbon Budget 2017](#)

Fossil fuel and cement emissions growth


The biggest changes in emissions were from a decline in coal and an increase in oil



Source: [CDIAC](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Carbon intensity of economic activity

Global emissions growth has generally recovered quickly from previous financial crises
It is unclear if the recent slowdown in global emissions is related to the Global Financial Crisis




Economic activity is measured in purchasing power parity (PPP) terms in 2010 US dollars.

Source: [CDIAC](#); [Peters et al 2012](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Emissions intensity per unit economic activity

The 10 largest economies have a wide range of emissions intensity of economic production




Emission intensity: CO₂ emissions from fossil fuel and industry divided by Gross Domestic Product

Source: [Global Carbon Budget 2017](#)

New generation of emissions scenarios

In the lead up to the IPCC's Sixth Assessment Report new scenarios have been developed to more systematically explore key uncertainties in future socioeconomic developments




Five Shared Socioeconomic Pathways (SSPs) have been developed to explore challenges to adaptation and mitigation. Shared Policy Assumptions (SPAs) are used to achieve target forcing levels (W/m²). Marker Scenarios are indicated.

Source: [Riahi et al. 2016](#); [IIASA SSP Database](#); [Global Carbon Budget 2017](#)

New generation of emissions scenarios

In the lead up to the IPCC's Sixth Assessment Report new scenarios have been developed to more systematically explore key uncertainties in future socioeconomic developments




Five Shared Socioeconomic Pathways (SSPs) have been developed to explore challenges to adaptation and mitigation. Shared Policy Assumptions (SPAs) are used to achieve target forcing levels (W/m²). Marker Scenarios are indicated.

Source: [Riahi et al. 2016](#); [IIASA SSP Database](#); [Global Carbon Budget 2017](#)

Pathways that avoid 2°C of warming


According to the Shared Socioeconomic Pathways (SSP) that avoid 2°C of warming, global CO₂ emissions need to decline rapidly and cross zero emissions after 2050



Source: [Riahi et al. 2016](#); [IIASA SSP Database](#); [Global Carbon Budget 2017](#)

CO₂ emissions and economic activity


In recent years, CO₂ emissions have been almost flat despite continued economic growth



Source: [Jackson et al 2017](#); [Global Carbon Budget 2017](#)

Kaya decomposition

The Kaya decomposition demonstrates the recent relative decoupling of economic growth from CO₂ emissions, driven by improved energy intensity




GWP: Gross World Product (economic activity), FFI: Fossil Fuel and Industry, Energy is Primary Energy from BP statistics using the substitution accounting method

Source: [Jackson et al 2017](#); [Global Carbon Budget 2017](#)

Emissions per capita

The 10 most populous countries span a wide range of development and emissions per person



Emission per capita: CO₂ emissions from fossil fuel and industry divided by population

Source: [Global Carbon Budget 2017](#)

Key statistics

Emissions 2016

Region/Country	Per capita tCO ₂ per person	Total		Growth 2015-16	
		GtCO ₂	%	GtCO ₂	%
Global (with bunkers)	4.8	36.18	100	0.163	0.0

OECD Countries

OECD	9.8	12.56	34.7	-0.110	-1.1
USA	16.5	5.31	14.7	-0.100	-2.1
OECD Europe	7.0	3.42	9.5	0.000	-0.3
Japan	9.5	1.21	3.3	-0.016	-1.6
South Korea	11.7	0.60	1.6	0.003	0.3
Canada	15.5	0.56	1.6	-0.005	-1.2

Non-OECD Countries

Non-OECD	3.6	22.25	61.5	0.220	0.7
China	7.2	10.15	28.1	0.000	-0.3
India	1.8	2.43	6.7	0.110	4.5
Russia	11.4	1.63	4.5	-0.036	-2.4
Iran	8.2	0.66	1.8	0.014	1.9
Saudi Arabia	19.7	0.63	1.8	0.011	1.4

International Bunkers

Aviation and Shipping	-	1.37	3.8	0.053	4.0
-----------------------	---	------	-----	-------	-----


Consumption-based Emissions

Consumption-based emissions allocate emissions to the location that goods and services are consumed

Consumption-based emissions = Production/Territorial-based emissions minus emissions embodied in exports plus the emissions embodied in imports

Consumption-based emissions (carbon footprint)


Allocating fossil and industry emissions to the consumption of products provides an alternative perspective.
USA and EU28 are net importers of embodied emissions, China and India are net exporters.



Consumption-based emissions are calculated by adjusting the standard production-based emissions to account for international trade
Source: [Peters et al 2011](#); [Le Quéré et al 2017](#); [Global Carbon Project 2017](#)

Consumption-based emissions (carbon footprint)

Transfers of emissions embodied in trade from non-Annex B countries to Annex B countries grew at over 11% per year between 1990 and 2007, but have since declined at over 1% per year.




Annex B countries were used in the Kyoto Protocol, but this distinction is less relevant in the Paris Agreement

Source: [CDIAC](#); [Peters et al 2011](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Major flows from production to consumption

Flows from location of generation of emissions to location of consumption of goods and services




Values for 2011. EU is treated as one region. Units: MtCO₂

Source: [Peters et al 2012](#)

Major flows from extraction to consumption

Flows from location of fossil fuel extraction to location of consumption of goods and services




Values for 2011. EU is treated as one region. Units: MtCO₂

Source: [Andrew et al 2013](#)

Land-use Change Emissions


Land-use change emissions

Land-use change emissions are highly uncertain. Higher emissions in 2016 are linked to increased fires during dry El Niño conditions in tropical Asia




Total global emissions

Total global emissions: 40.8 ± 2.7 GtCO₂ in 2016, 52% over 1990
 Percentage land-use change: 42% in 1960, 12% averaged 2007-2016



Total global emissions by source

Land-use change was the dominant source of annual CO₂ emissions until around 1950




Others: Emissions from cement production and gas flaring

Source: [CDIAC](#); [Houghton and Nassikas 2017](#); [Hansis et al 2015](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Historical cumulative emissions by source

Land-use change represents about 31% of cumulative emissions over 1870–2016, coal 32%, oil 25%, gas 10%, and others 3%



Others: Emissions from cement production and gas flaring

Source: [CDIAC](#); [Houghton and Nassikas 2017](#); [Hansis et al 2015](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Closing the Global Carbon Budget

Fate of anthropogenic CO₂ emissions (2007–2016)

Sources = Sinks



34.3 GtCO₂/yr
88%



12%
4.9 GtCO₂/yr

17.3 GtCO₂/yr
47%



30%
11.2 GtCO₂/yr



23%
8.7 GtCO₂/yr



Budget Imbalance:


(the difference between estimated sources & sinks)

6%

2.1 GtCO₂/yr


Global carbon budget

Carbon emissions are partitioned among the atmosphere and carbon sinks on land and in the ocean
The “imbalance” between total emissions and total sinks reflects the gap in our understanding




Changes in the budget over time

The sinks have continued to grow with increasing emissions, but climate change will affect carbon cycle processes in a way that will exacerbate the increase of CO₂ in the atmosphere




Global Carbon Project



The budget imbalance is the total emissions minus the estimated growth in the atmosphere, land and ocean.
It reflects the limits of our understanding of the carbon cycle.


Source: [CDIAC](#); [NOAA-ESRL](#); [Houghton and Nassikas 2017](#); [Hansis et al 2015](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

The atmospheric concentration growth rate has shown a steady increase
The high growth in 1987, 1998, & 2015-16 reflect a strong El Niño, which weakens the land sink



Source: [NOAA-ESRL](#); [Global Carbon Budget 2017](#)


The ocean carbon sink continues to increase
 $8.7 \pm 2 \text{ GtCO}_2/\text{yr}$ for 2007–2016 and $9.6 \pm 2 \text{ GtCO}_2/\text{yr}$ in 2016



Source: [SOCATv5](#); [Bakker et al 2016](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)


Individual estimates from: Aumont and Bopp (2006); Buitenhuis et al. (2010); Doney et al. (2009); Hauck et al. (2016); Ilyina et al. (2013); Landschützer et al. (2016); Law et al. (2017); ; Rödenbeck et al. (2014). Séférian et al. (2013); Schwinger et al. (2016). Full references provided in Le Quéré et al. (2017).

The land sink was 11.2 ± 3 GtCO₂/yr during 2007-2016 and 10 ± 3 GtCO₂/yr in 2016
Total CO₂ fluxes on land (including land-use change) are constrained by atmospheric inversions



Total land and ocean fluxes

Total land and ocean fluxes show more interannual variability in the tropics




Source: [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

Individual estimates from: Aumont and Bopp (2006); Buitenhuis et al. (2010); Chevallier et al. (2005); Clarke et al. (2011); Doney et al. (2009); Guimberteau et al. (2017); Hauck et al. (2016); Haverd et al. (2017); Ilyina et al. (2013); Jain et al. (2013); Kato et al. (2013); Keller et al. (2017); Krinner et al. (2005); Landschützer et al. (2016); Law et al. (2017); Melton and Arora (2016); Oleson et al. (2013); Reick et al. (2013); Rödenbeck et al. (2003); Rödenbeck et al. (2014); Séférian et al. (2013); Schwinger et al. (2016); Sitch et al. (2003); Smith et al. (2014); Tian et al. (2015); van der Laan-Luijkx et al. (2017); Woodward et al. (1995); Zaehle and Friend (2010). Full references provided in Le Quéré et al. (2017).

Remaining carbon budget imbalance

Large and unexplained variability in the global carbon balance caused by uncertainty and understanding hinder independent verification of reported CO₂ emissions



The budget imbalance is the carbon left after adding independent estimates for total emissions, minus the atmospheric growth rate and estimates for the land and ocean carbon sinks using models constrained by observations

Source: [Le Quéré et al 2017](#); [Global Carbon Budget 2017](#)

The cumulative contributions to the global carbon budget from 1870
The carbon imbalance represents the gap in our current understanding of sources and sinks





Figure concept from [Shrink That Footprint](#)

Source: [CDIAC](#); [NOAA-ESRL](#); [Houghton and Nassikas 2017](#); [Hansis et al 2015](#); [Joos et al 2013](#); [Khatiwala et al. 2013](#); [DeVries 2014](#); [Le Quéré et al 2017](#); [Global Carbon Budget 2016](#)


Atmospheric concentration

The global CO₂ concentration increased from ~277ppm in 1750 to 403ppm in 2016 (up 45%)
2016 was the first full year with concentration above 400ppm



Trends in CO₂ emissions and concentrations


Atmospheric CO₂ concentration had record growth in 2015 & 2016 due to record high emissions and El Niño conditions, but growth is expected to reduce due to the end of El Niño



Source: [Peters et al 2017](#); [Global Carbon Budget 2017](#)

Verification of a sustained change in CO₂ emissions

Our ability to detect changes in CO₂ emissions based on atmospheric observations is limited by our understanding of carbon cycle variability




Observations show a large-interannual to decadal variability, which can only be partially *reconstructed* through the global carbon budget. The difference between observations and reconstructed is the “budget imbalance”.

Source: [Peters et al 2017](#); [Global Carbon Budget 2017](#)

Seasonal variation of atmospheric CO₂ concentration

Weekly CO₂ concentration measured at Mauna Loa stayed above 400ppm throughout 2016 and is forecast to average 406.8 in 2017



Forecasts are [an update](#) of [Betts et al 2016](#). The deviation from monthly observations is 0.24 ppm (RMSE).

Updates of [this figure](#) are available, and [another](#) on the drivers of the atmospheric growth

Source: Tans and Keeling (2017), [NOAA-ESRL](#), [Scripps Institution of Oceanography](#)

End notes



Global Carbon Budget 2017

In 2017, CO₂ emissions from fossil fuels and industry are projected to **grow by 2.0%** (+0.8 to +3.0%). This follows three years of nearly **no growth (2014-2016)**

In 2016 atmospheric CO₂ levels reached **403 ppm**...
...and are projected to increase by 2.5 ppm in 2017 (+2.0 to +3.0ppm)


315 ppm
Data: Scripps/NOAA-ESRL
1960 2016

The **plateau** of last year was not peak emissions after all...



...we are changing trajectory...

Emissions **decreased** significantly in the presence of a growing GDP in **22 countries (representing 20% of global emissions)** in the last decade (2007-2016). Other **notable changes** are also shown



Brazil: Emissions declining but probably due to economic crisis


China: Emissions declined for the past 3 years but are up again

Japan: Emissions declined recently


India: Emissions grew 6% in the past decade but slowed in 2017

...but atmospheric concentrations continue to rise

The carbon cycle has both **emissions sources** and **carbon sinks**, and their difference is the **atmospheric growth** (2007-2016)



Atmospheric growth increases in line with **total CO₂ emissions**, but has large variability. The **2015-2016 El Niño** led to a record high growth due to lower CO₂ uptake by tropical forests



Copyright: CC BY-NC-SA

Produced by the Future Earth Media Lab for the Global Carbon Project: <http://www.globalcarbonproject.org/carbonbudget/index.htm>
Written and edited by Peter Canadell (CSIRO), Robbie Andrew and Glen Peters (CI CERF), and Corinne Le Quéré (Tyndall Centre UEA)
with the Global Carbon Budget team. Updated from 2016 version. Infographic by Nigel Hawton.
Credits: Le Quéré et al. Earth System Science Data Discussions (2017), NOAA-ESRL and the Scripps Institution of Oceanography, CDIAC/NDC projection based on UNFCCC analysis based of Rogelj et al Nature 2016 assuming constant CO₂/GHG ratio

The work presented in the **Global Carbon Budget 2017** has been possible thanks to the contributions of **hundreds of people** involved in observational networks, modeling, and synthesis efforts.

We thank the institutions and agencies that provide support for individuals and funding that enable the collaborative effort of bringing all components together in the carbon budget effort.

We thank the sponsors of the GCP and GCP support and liaison offices.

We also want to thank each of the many funding agencies that supported the individual components of this release. A full list is provided in Table B1 of Le Quéré et al. 2017.

<https://doi.org/10.5194/essdd-2017-123>

We also thank the Fondation BNP Paribas for supporting the Global Carbon Atlas.

This presentation was created by Robbie Andrew with Pep Canadell, Glen Peters and Corinne Le Quéré in support of the international carbon research community.

futureearth
research for global sustainability



The Research Council
of Norway



European
Commission

°CICERO
Center for International
Climate and Environmental
Research - Oslo



UEA
University of East Anglia

Tyndall°Centre®
for Climate Change Research



FONDATION
BNP PARIBAS



Global Carbon Project

Attribution 4.0 International (CC BY 4.0)

This deed highlights only some of the key features and terms of the actual license. It is not a license and has no legal value. You should carefully review all of the terms and conditions of the actual license before using the licensed material. This is a human-readable summary of (and not a substitute for) the [license](#).

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions — You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.

You do not have to comply with the license for elements of the material in the public domain or where your use is permitted by an applicable exception or limitation. No warranties are given. The license may not give you all of the permissions necessary for your intended use. For example, other rights such as publicity, privacy, or moral rights may limit how you use the material.